

Durham Research Online

Deposited in DRO:

08 February 2021

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Straughan, B. (2021) 'Continuous dependence for the Brinkman–Darcy–Kelvin–Voigt equations backward in time.', *Mathematical methods in the applied sciences.*, 44 (6). pp. 4999-5004.

Further information on publisher's website:

<https://doi.org/10.1002/mma.7082>

Publisher's copyright statement:

This is the peer reviewed version of the following article: Straughan, B. (2021). Continuous dependence for the Brinkman–Darcy–Kelvin–Voigt equations backward in time. *Mathematical Methods in the Applied Sciences* 44(6): 4999-5004., which has been published in final form at <https://doi.org/10.1002/mma.7082>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Continuous dependence for the Brinkman - Darcy - Kelvin - Voigt equations backward in time

B. Straughan
Department of Mathematics
University of Durham, Durham DH1 3LE, UK

August 19, 2020

Abstract

We show that the solution to the Brinkman - Darcy - Kelvin - Voigt equations backward in time depends Hölder continuously upon the final data. A logarithmic convexity technique is employed and uniqueness of the solution is simultaneously achieved.

1 Introduction

The problem of analysing the solution to an improperly posed problem for a system of partial differential equations has attracted many writers over the years and continues to do so recently, see e.g. Agmon [1966], Agmon and Nirenberg [1967], Ames and Hughes [2005], Ames and Straughan [1997], Benrabah et al. [2020], Caffisch et al. [2017], Carasso [2013, 2019, 2020], Chirita [2014], Chirita and Zampoli [2015], Fury [2020], Fury and Hughes [2012], Hetrick and Hughes [2009], John [1960], Knops and Payne [1968], Yang and Deng [2017]cw., In particular, the pioneering paper of John [1960] showed how one could recover a restricted class of stable solutions by requiring an *a priori* bound at one particular place. This work has influenced many of the subsequent articles.

There has been a considerable amount of work dealing with solutions to the Navier - Stokes equations backward in time with regard to establishing uniqueness, stability (in a sense like that of John [1960]), and structural stability, cf. Ames and Payne [1994], Carasso [2020], Crispo et al. [2019], Galdi and Straughan [1988], Harfash [2013, 2014], Payne [1971, 1992, 1993, 1975], Payne and Straughan [1989, 1999], Straughan [1983]. Such problems are of practical value in extrapolating from the past and computational methods may be based on analytical results, as explained in detail by Carasso [2020].

In addition to the Navier - Stokes equations for flow of a linearly viscous incompressible fluid, there has been considerable interest in describing fluids which are viscous but remember some of their past history, so called viscoelastic fluids, see e.g. Amendola and Fabrizio [2010], Amendola et al. [2009], Fabrizio

et al. [2015], Franchi et al. [2014, 2015a,b], Gatti et al. [2005]. A special class of these viscoelastic models is known as Kelvin - Voigt materials, Chirita and Zampoli [2015], and applied as extensions of the Navier - Stokes theory through the Navier - Stokes - Voigt equations, see e.g. Berselli and Bisconti [2012], Di Plinio et al. [2018], Sviridyuk and Sukacheva [1998].

The goal of this work is to establish an appropriate stability estimate for the solution to the Navier - Stokes - Voigt equations backward in time, which we do by implementing a suitable logarithmic convexity method. Such techniques are described in a variety of contexts by e.g. Agmon [1966], Ames and Straughan [1997], Carasso [1994, 1999, 2013, 2019, 2020], Crispo et al. [2019], Knops and Payne [1968], Payne [1975].

2 The Navier - Stokes -Voigt equations

Let $\Omega \subset \mathbb{R}^3$ be a bounded domain with boundary Γ sufficiently smooth to allow use of the divergence theorem. The inner product and norm on $L^2(\Omega)$ will be denoted by (\cdot, \cdot) and $\|\cdot\|$, respectively.

If the velocity field is denoted by v_i then the Navier - Stokes - Voigt equations may be written as

$$\begin{aligned} v_{i,t} - \alpha \Delta v_{i,t} + v_j v_{i,j} &= \Delta v_i - p_{,i}, \\ v_{i,i} &= 0, \end{aligned} \tag{1}$$

where without loss of generality for the analysis herein the viscosity and density have been taken to have value 1, $\alpha > 0$ is a constant, p is the pressure field, Δ is the Laplacian, standard indicial notation is employed in conjunction with the Einstein summation convention, and $,_t$ denotes $\partial/\partial t$, whereas $,_i$ denotes $\partial/\partial x_i$.

In this work we reverse time to analyse a solution to the boundary - initial value problem backward in time. Di Plinio et al. [2018] allow (1) to also contain a Rayleigh friction term (linear in v_i) and we include this. The relevant equations backward in time may then be written

$$\begin{aligned} v_{i,t} - \alpha \Delta v_{i,t} &= v_j v_{i,j} - \Delta v_i + p_{,i} + \beta v_i, \\ v_{i,i} &= 0, \end{aligned} \tag{2}$$

where β is a constant. Equations (2) hold on the domain $\Omega \times (0, T]$, for $T < \infty$, and the boundary - initial value problem is completed with boundary conditions

$$v_i(\mathbf{x}, t) = h_i(\mathbf{x}, t), \quad \mathbf{x} \in \Gamma, \quad t > 0, \tag{3}$$

and the initial conditions

$$v_i(\mathbf{x}, 0) = g_i(\mathbf{x}), \quad \mathbf{x} \in \Omega. \tag{4}$$

The boundary - initial value problem consisting of (2) - (4) is denoted by \mathcal{P} .

3 Stability backward in time

To study stability of a solution to \mathcal{P} we let u_i, p and v_i, q be two solutions for the same boundary data h_i , but for different initial data functions g_i^u and g_i^v . Define $w_i = u_i - v_i$ and $\pi = p - q$, and then from (2) - (4) one finds w_i satisfies the boundary - initial value problem

$$\begin{aligned} w_{i,t} - \alpha \Delta w_{i,t} &= u_j w_{i,j} + w_j v_{i,j} - \Delta w_i + \pi_{,i} + \beta w_i, \\ w_{i,i} &= 0, \end{aligned} \quad (5)$$

holding on $\Omega \times (0, T]$, with boundary conditions

$$w_i(\mathbf{x}, t) = 0, \quad \mathbf{x} \in \Gamma, \quad t > 0, \quad (6)$$

and initial conditions

$$w_i(\mathbf{x}, 0) = w_i^0(\mathbf{x}), \quad \mathbf{x} \in \Omega, \quad (7)$$

where $w_i^0 = g_i^u - g_i^v$.

Let $\|\cdot\|_p$ denote the norm on $L^p(\Omega)$. We now introduce the constraint classes for a function. We say $\phi_i \in \mathcal{M}_1$ if

$$\sup_{\Omega \times (0, T]} \phi_i \phi_i \leq M_1^2, \quad (8)$$

and $\psi_i \in \mathcal{M}_2$ if

$$\sup_{\Omega \times (0, T]} \{ \|(\psi_{i,j} - \psi_{j,i})(\psi_{i,j} - \psi_{j,i})\|_{3/2} + \|\psi_{i,t} \psi_{i,t}\|_{3/2} \} \leq M_2^2, \quad (9)$$

for constants M_1 and M_2 .

Theorem 1

Suppose u_i, v_i satisfy \mathcal{P} as outlined above with $u_i \in \mathcal{M}_1$, $v_i \in \mathcal{M}_2$. Let also M be an a priori bound for w_i at time T in the following sense,

$$\|\mathbf{w}(T)\|^2 + \alpha \|\nabla \mathbf{w}(T)\|^2 \leq M < \infty. \quad (10)$$

Then the solution to \mathcal{P} depends Hölder continuously upon the initial data on compact subsets of $[0, T)$.

Proof. Define the function $F(t)$ by

$$F(t) = \|\mathbf{w}(t)\|^2 + \alpha \|\nabla \mathbf{w}(t)\|^2. \quad (11)$$

By differentiation,

$$F'(t) = 2(w_i, w_{i,t}) + 2\alpha(w_{i,j}, w_{i,jt}). \quad (12)$$

Next, integrate by parts the $w_{i,jt}$ term and then employ equations (5), to find after further integration by parts and use of the boundary conditions

$$F'(t) = 2\|\nabla \mathbf{w}\|^2 + 2(w_i, w_j v_{i,j}) + 2\beta \|\mathbf{w}\|^2.$$

Next differentiate again to obtain after some integration by parts

$$F''(t) = 2(w_{i,t}, w_j v_{i,j}) + 2(w_i, w_{j,t} v_{i,j}) + 2(w_i, w_j v_{i,jt}) \\ - 4(w_{i,t}, \Delta w_i) + 4\beta(w_i, w_{i,t}).$$

One now substitutes for Δw_i from (2)₁ and after further integration by parts and use of the boundary conditions one may arrive at the expression

$$F''(t) = 4\|\dot{\mathbf{w}}\|^2 + 4\alpha\|\nabla \dot{\mathbf{w}}\|^2 - 4(w_{i,t}, u_j w_{i,j}) \\ - 2(w_{i,t}, w_j[v_{i,j} - v_{j,i}]) - 2(w_{i,j}, w_j v_{i,t}), \quad (13)$$

where $\dot{\mathbf{w}} \equiv w_{i,t}$. Introduce the function χ_i by

$$\chi_i = w_{i,t} - \frac{1}{2}u_j w_{i,j} - \frac{1}{4}w_j(v_{i,j} - v_{j,i}).$$

Then F'' in (13) may be rewritten in the form

$$F''(t) = 4\|\chi\|^2 + 4\alpha\|\nabla \dot{\mathbf{w}}\|^2 - 2(w_j, w_{i,j} v_{i,t}) - \|\mathbf{u} \cdot \nabla \mathbf{w}\|^2 \\ - (u_j w_{i,j}, w_k[v_{i,k} - v_{k,i}]) - \frac{1}{4}\|w_j(v_{i,j} - v_{j,i})\|^2, \quad (14)$$

where

$$\|w_j(v_{i,j} - v_{j,i})\|^2 = \int_{\Omega} w_j(v_{i,j} - v_{j,i}) w_k(v_{i,k} - v_{k,i}) dx.$$

Upon employing the arithmetic - geometric mean and Cauchy - Schwarz inequalities on the right of (14) we may arrive at

$$F''(t) \geq 4\|\chi\|^2 + 4\alpha\|\nabla \dot{\mathbf{w}}\|^2 - 2(w_j, w_{i,j} v_{i,t}) \\ - 2\|\mathbf{u} \cdot \nabla \mathbf{w}\|^2 - \frac{1}{2}\|w_j(v_{i,j} - v_{j,i})\|^2. \quad (15)$$

We next estimate the last three terms on the right of (15).

Firstly use the Cauchy - Schwarz and Hölder inequalities to see that

$$(w_j, w_{i,j} v_{i,t}) \leq \frac{\epsilon}{2} \int_{\Omega} |\mathbf{w}|^2 |\dot{\mathbf{v}}|^2 dx + \frac{1}{2\epsilon} \|\nabla \mathbf{w}\|^2 \\ \leq \frac{\epsilon}{2} \|\mathbf{w}\|_6^2 \left(\int_{\Omega} |\dot{\mathbf{v}}|^3 dx \right)^{2/3} + \frac{1}{2\epsilon} \|\nabla \mathbf{w}\|^2.$$

Now use the Sobolev inequality $\|\mathbf{w}\|_6 \leq c\|\mathbf{w}\|_{H^1}$ for c constant, and then select $\epsilon = 1/M_2 c$ where M_2 is defined in (9). In this way we derive

$$2(w_j, w_{i,j} v_{i,t}) \leq 2M_2 c \|\nabla \mathbf{w}\|^2. \quad (16)$$

Using (8) we deduce

$$2\|\mathbf{u} \cdot \nabla \mathbf{w}\|^2 \leq 2M_1^2 \|\nabla \mathbf{w}\|^2. \quad (17)$$

Finally, using (9) and the Sobolev inequality,

$$\begin{aligned} \frac{1}{2} \|w_j(v_{i,j} - v_{j,i})\|^2 &\leq \frac{1}{2} \|\mathbf{w}\|_6^2 \left(\int_{\Omega} [(v_{i,j} - v_{j,i})(v_{i,j} - v_{j,i})]^{3/2} dx \right)^{2/3} \\ &\leq \frac{1}{2} c^2 M_2^2 \|\nabla \mathbf{w}\|^2. \end{aligned} \quad (18)$$

Employing (16) - (18) we find that the last three terms on the right of (15) may be bounded below by the term $-k_1 \|\nabla \mathbf{w}\|^2$ where

$$k_1 = 2M_2c + \frac{1}{2}c^2M_2^2 + 2M_1^2.$$

Thus, from (15) we find

$$\begin{aligned} F''(t) &\geq 4\|\chi\|^2 + 4\alpha\|\nabla \dot{\mathbf{w}}\|^2 - k_1\|\nabla \mathbf{w}\|^2 \\ &\geq 4\|\chi\|^2 + 4\alpha\|\nabla \dot{\mathbf{w}}\|^2 - kF, \end{aligned} \quad (19)$$

where $k = k_1/\alpha$.

We now form the combination $FF'' - (F')^2$ using (19), (12) and (11), and note that $(\chi_i, w_i) = (w_i, w_{i,t})$. Then from (19) we may establish the inequality

$$FF'' - (F')^2 \geq 4S^2 - kF^2, \quad (20)$$

where

$$\begin{aligned} S^2 &= (\|\mathbf{w}\|^2 + \alpha\|\nabla \mathbf{w}\|^2)(\|\chi\|^2 + \alpha\|\nabla \dot{\mathbf{w}}\|^2) \\ &\quad - [(w_i, \chi_i) + \alpha((\nabla \mathbf{w}, \nabla \dot{\mathbf{w}}))]^2, \end{aligned}$$

which is non-negative by virtue of the Cauchy - Schwarz inequality. Whence from (20) we obtain

$$FF'' - (F')^2 \geq -kF^2. \quad (21)$$

Upon dividing by F^2 we may see that

$$\frac{d^2}{dt^2} \left[\log \{ F \exp(kt^2/2) \} \right] \geq 0. \quad (22)$$

Using the properties of a convex function one may then show that

$$F(t) \leq \exp \left[\frac{kt}{2}(T-t) \right] [F(0)]^{(T-t)/T} [F(T)]^{t/(T-t)}. \quad (23)$$

The *a priori* bound of the theorem, (10), is now utilized and we derive from (23)

$$F(t) \leq K [F(0)]^{(T-t)/T}, \quad t \in [0, T], \quad (24)$$

where $K(t) = M^{t/(T-t)} \exp[kt(T-t)/2]$. The inequality (24) establishes Hölder continuity for a solution to \mathcal{P} on a compact sub-interval of $[0, T]$ for a solution in the measure $\|\mathbf{w}\|$ or $\|\nabla \mathbf{w}\|$ and the theorem is proved.

Corollary 1

Under the conditions of theorem 1 the solution to \mathcal{P} is unique.

Proof. By contradiction. Suppose u_i is not identically zero and $u_i \neq 0$ for $t > 0$ whence $F(t) > 0$ for $t \in [\epsilon, T]$. In this case we divide (21) by F^2 which leads to (22) on $[\epsilon, T]$. Fix $t \in (\epsilon, T)$ and from (22) we know by properties of a convex function,

$$\begin{aligned} -\infty < \log \left[F(t) \exp\left(\frac{kt^2}{2}\right) \right] &\leq \left(\frac{T-t}{T-\epsilon}\right) \log \left[F(\epsilon) \exp\left(\frac{k\epsilon^2}{2}\right) \right] \\ &\quad + \left(\frac{t-\epsilon}{T-\epsilon}\right) \log \left[F(T) \exp\left(\frac{kT^2}{2}\right) \right]. \end{aligned}$$

For uniqueness $F(0) = 0$, so we let $\epsilon \rightarrow 0$ to obtain a contradiction since t is fixed, and the proof of the Corollary follows.

4 Conclusions

We have derived a precise estimate for a solution to the Navier - Stokes - Voigt equations backward in time to depend Hölder continuously on the initial data on compact subsets of the interval $[0, T)$. We also allowed for the generalized Navier - Stokes - Voigt system which includes the Rayleigh friction term suggested in the work of Di Plinio et al. [2018]. The estimate obtained in (24) could be combined with the ideas of Carasso [2020] to allow one to compute accurate solutions to the Navier - Stokes - Voigt equations backward in time.

References

- S. Agmon. Unicité et convexité dans les problèmes différentiels. In *Sem. Math. Sup.* University of Montreal Press, 1966.
- S. Agmon and L. Nirenberg. Lower bounds and uniqueness theorems for solutions of differential equations in a Hilbert space. *Comm. Pure Appl. Math.*, 20:207–229, 1967.
- K. A. Ames and R. J. Hughes. Structural stability for ill - posed problems in Banach space. *Semigroup Forum*, 85:127–145, 2005.
- K. A. Ames and B. Straughan. *Non - standard and improperly posed problems*. Academic Press, 1997.
- A. Benrabah, N. Boussetila, and F. Rebbani. Modified auxilliary boundary conditions method for an ill - posed problem for the homogeneous biharmonic equation. *Math. Meth. Appl. Sci.*, 43:358–383, 2020.
- R. E. Caffisch, F. Gargano, and V. Sammartino, M. and Sciacca. Regularized Euler - α motion of an infinite array of vortex sheets. *Boll. Unione Matematica Italiana*, 10:113–141, 2017.

- A. Carasso. Reconstructing the past from imprecise knowledge of the present: some examples of non uniqueness in solving parabolic equations backward in time. *Math. Meth. Appl. Sci.*, 36:249–261, 2013.
- A. Carasso. Stable explicit stepwise marching scheme in ill posed time - reversed 2D Burgers' equation. *Inverse Prob. Sci. Engng.*, 27:1672–1688, 2019.
- A. Carasso. Computing ill posed time - reversed 2D Navier - Stokes equations using a stabilized explicit finite difference scheme marching backward in time. *Inverse Prob. Sci. Engng.*, 28:doi:10.1080/17415977.2019.1698564, 2020.
- S. Chirita. Well - posed problems. In R. B. Hetnarski, editor, *Encyclopedia of thermal stresses*. Springer, Dordrecht, 2014. https://doi.org/10.1007/978-94-007-2739-7_264.
- S. Chirita and V. Zampoli. On the forward and backward in time problems in the Kelvin - Voigt thermoelastic materials. *Mech. Res. Comm.*, 68:25–30, 2015.
- M. A. Fury. Ill - posed problems associated with sectorial operators in Banach space. *J. Math. Anal. Appl.*, 400:<https://doi.org/10.1016/j.jmaa.2020.124484>, 2020.
- M. A. Fury and R. J. Hughes. Regularization for a class of ill - posed problems in Banach space. *Semigroup Forum*, 85:191–212, 2012.
- B. M. C. Hetrick and R. J. Hughes. Continuous dependence on modelling for nonlinear ill - posed problems. *J. Math. Anal. Appl.*, 349:420–435, 2009.
- F. John. Continuous dependence on data for solutions of partial differential equations with a prescribed bound. *Comm. Pure Appl. Math.*, 13:551–585, 1960.
- R. J. Knops and L. E. Payne. On the stability of solutions of the Navier - Stokes equations backward in time. *Arch. Rational Mech. Anal.*, 29:331–335, 1968.
- X. M. Yang and Z. L. Deng. A data assimilation process for linear ill - posed problems. *Math. Meth. Appl. Sci.*, 40:5831–5840, 2017.
- K. A. Ames and L. E. Payne. Continuous dependence results for solutions of the Navier - Stokes equations backward in time. *Nonlinear Anal., Theory, Meths., Applicns.*, 23:103–113, 1994.
- F. Crispo, C. R. Grisanti, and P. Maremonti. Some new properties of a suitable weak solution to the Navier - Stokes equations. [arXiv:1904.07641v2](https://arxiv.org/abs/1904.07641v2), 2019.
- G. P. Galdi and B. Straughan. Stability of solutions of the Navier - Stokes equations backward in time. *Arch. Rational Mech. Anal.*, 101:107–114, 1988.

- A. J. Harfash. Continuous dependence on the coefficients for double diffusive convection in Darcy flow with magnetic field effect. *Anal. Math. Phys.*, 3: 163–181, 2013.
- A. J. Harfash. Structural stability for two convection models in a reacting fluid with magnetic field effect. *Ann. Henri Poincaré*, 15:2441–2465, 2014.
- L. E. Payne. Uniqueness and continuous dependence criteria for the Navier - Stokes equations. *Rocky Mount. J. Math.*, 2:641–660, 1971.
- L. E. Payne. Some remarks on ill - posed problems for viscous fluids. *Int. J. Engng. Sci.*, 30:1341–1347, 1992.
- L. E. Payne. Continuous dependence on spatial geometry for solutions of the Navier - Stokes equations backward in time. *Nonlinear Anal., Theory, Meths., Applicns.*, 21:651–664, 1993.
- L. E. Payne. *Improperly posed problems in partial differential equations*. SIAM, 1975. Regional Conf. Ser., Appl. Math.
- L. E. Payne and B. Straughan. Comparison of viscous flows backward in time with small data. *Int. J. Non-Linear Mech.*, 24:209–214, 1989.
- L. E. Payne and B. Straughan. Effect of errors in the spatial geometry for temperature dependent Stokes flow. *J. Math. Pures Appl.*, 78:609–632, 1999.
- B. Straughan. Backward uniqueness and unique continuation for solutions to the Navier - Stokes equations on an exterior domain. *J. Math. Pures Appl.*, 62:49–62, 1983.
- G. Amendola and M. Fabrizio. Thermal convection in a simple fluid with fading memory. *J. Math. Anal. Appl.*, 366:444–459, 2010.
- G. Amendola, M. Fabrizio, M. Golden, and B. Lazzari. Free energies and asymptotic behaviour for incompressible viscoelastic fluids. *Applicable Anal.*, 88: 789–805, 2009.
- M. Fabrizio, B. Lazzari, and R. Nibbi. Aymptotic stability in linear viscoelasticity with supplies. *J. Math. Anal. Appl.*, 427:629–645, 2015.
- F. Franchi, B. Lazzari, and R. Nibbi. Uniqueness and stability results for non-linear Johnson - Segalman viscoelasticity and related models. *Disc. Cont. Dyn. Sys. B*, 19:2111–2132, 2014.
- F. Franchi, B. Lazzari, and R. Nibbi. Mathematical models for the non - isothermal Johnson - Segalman viscoelasticity in porous media: stability and wave propagation. *Math. Meth. Appl. Sci.*, 38:4075–4087, 2015a.
- F. Franchi, B. Lazzari, and R. Nibbi. The Johnson - Segalman model versus a non - ideal MHD theory. *Phys. Lett. A*, 379:1431–1436, 2015b.

- S. Gatti, C. Giorgi, and V. Pata. Navier - Stokes limit of Jeffreys type flows. *Physica D*, 203:55–79, 2005.
- L. C. Berselli and L. Bisconti. On the structural stability of the Euler - Voigt and Navier - Stokes - Voigt models. *Nonlinear Analysis*, 75:117–130, 2012.
- F. Di Plinio, A. Giorgini, V. Pata, and R. Temam. Navier - Stokes - Voigt equations with memory in 3D lacking instantaneous kinematic viscosity. *J. Nonlinear Sci*, 28:653–686, 2018.
- G. A. Sviridyuk and T. G. Sukacheva. On the solvability of a nonstationary problem describing the dynamics of an incompressible viscoelastic fluid. *Mathematical Notes*, 63:388–395, 1998.
- A. Carasso. Overcoming Hölder continuity in ill posed continuation problems. *SIAM J. Numer. Anal.*, 31:1535–1557, 1994.
- A. Carasso. Logarithmic convexity and the “slow evolution” constraint in ill posed initial value problems. *SIAM J. Math. Anal.*, 30:479–496, 1999.